

FEMALE SEXUAL FUNCTION AND URINARY INCONTINENCE AFTER HIFEM[®] PROCEDURE

FEMALE SEXUAL FUNCTION AND URINARY INCONTINENCE: SIX MONTH FOLLOW-UP AFTER HIGH-INTENSITY FOCUSED ELECTROMAGNETIC PROCEDURE


Julene B. Samuels MD, FACS¹, Kimberly Evans MD, FACOG²

1. Julene B Samuels MD, FACS, Prospect, KY, USA
2. Hillcroft Medical Clinic, Sugar Land, TX, USA

Presented at North American Menopause Society 2020 Virtual Annual Meeting

HIGHLIGHTS

- **31 females** (49.8±9.9 years), who showed a decreased interest in sexual activity accompanied by UI, were enrolled in this prospective one-arm study.
- Subjects' **continence** and **sexual function** have been significantly ($P<0.001$) **improved**.
- Greatest improvement was seen after 3 months and it was maintained for 6 months.
 - **ICIQ-SF** overall score decreased on average by **61.6%** (-7.0 points)
 - **FSFI** showed significant increase by **35.5%** (+7.9 points)
 - **PISQ-12** score increased significantly by **26.3%** (+8.4 points)
- Patients benefited from increased sexual desire, improved lubrication, and arousal, and they were also more satisfied with their orgasm intensity.



ICIQ-SF (left) and FSFI (right) scores were significantly ($P<0.001$) improved after the HIFEM treatments.
Red dotted line indicates the threshold for female sexual dysfunction equal to 26.55 points.